

Current level of influenza activity: *Baseline activity (influenza not circulating)*

Influenza activity trend: *Stable*

Confirmed influenza cases since 2020 week 40: 36 (nine influenza A(H3N2), 14 influenza A(not subtyped) and 13 influenza B).

Key points – Wales


Surveillance indicators suggest that influenza is not currently circulating in Wales.

The sentinel GP consultation rate for influenza-like illness (ILI) increased during week 22 (ending 06/06/2021). During week 22, no cases of influenza were confirmed. COVID-19 cases continue to be detected in symptomatic patients in hospital and in the community, although levels are low compared to the 2020-21 winter. Rhinovirus remains the most commonly detected cause of non-COVID-19 Acute Respiratory Infection (ARI) and has increased from Week 10 onwards, as have a number of other causes of ARI. Surveillance data suggest that influenza and Respiratory Syncytial Virus (RSV) have not circulated over the winter months, but there remains potential for later unseasonal activity. There has been a greater diversity of causes of ARI in recent weeks with notable increases in adenovirus and parainfluenza.

- The **Sentinel GP consultation rate for influenza-like illness (ILI)** in Wales during week 22 was 0.5 consultations per 100,000 practice population (Table 1). This increased compared to the previous week (0.0 consultations per 100,000) but remains well below baseline threshold for seasonal influenza activity (11.0 per 100,000 practice population) (Figure 1). Caution should be used when comparing consultation rates from March 2020 onwards to previous periods due to the changes in health-seeking behaviours brought about by the COVID-19 pandemic.
- The **Sentinel GP consultation rate for Acute Respiratory Infections (ARI)** was 64.1 per 100,000 practice population during week 22, this is a slight increase compared to the previous week (63.8 per 100,000) (Table 2). Weekly consultations for Upper Respiratory Tract Infections decreased and Lower Respiratory Tract Infections increased compared to the previous week. The age-group specific consultation rate for ARI during week 22 was highest in one to four year olds (455.4 per 100,000 practice population).
- The percentage of calls to **NHS Direct Wales** which were 'influenza-related' (cold/flu, cough, fever, headache and sore throat) during week 22 increased to 15.5% (Figure 8).
- During week 22, 1,314 specimens received respiratory panel testing from hospital and non-sentinel GP patients with ARI. These results do not include samples tested solely for SARS-CoV2. There were 198 rhinoviruses, 49 adenoviruses, 33 parainfluenza, nine enteroviruses and eight seasonal coronaviruses detected in week 22 (Figure 4). Additionally, 3,241 samples from patients were tested for influenza, RSV and SARS-CoV2 only, a proportion of these tests may be associated with screening activities rather than diagnostic testing for patients presenting with ARI. Of these 3,241 samples, none were positive for influenza, two were positive for RSV and 27 were positive for SARS-CoV2 (figure 5). Forty-four respiratory specimens were tested from patients in intensive care units (ICU) and none were positive for influenza (Figure 6). For the latest COVID-19/ SARS-CoV2 surveillance data please see the [PHW daily dashboard](#)
- No surveillance samples from patients with ILI, collected by **sentinel GPs** during week 22, had been received by Public Health Wales Microbiology as at 09/06/2021.
- There has been no detectable RSV season in 2020-21. A small number of cases have been confirmed in recent weeks, but activity remains well below baseline.
- During week 22, **16 ARI outbreaks** were reported to the Public Health Wales Health Protection team, all were reported as COVID-19 outbreaks. Seven were in residential homes, two were in a school/nursery setting and seven were in a community, mixed or other setting.
- According to [EuroMoMo](#) analysis, all-cause deaths were not significantly in excess during week 20 (latest data).


Respiratory infection activity in Wales

Figure 1. Clinical consultation rate for ILI per 100,000 practice population in Welsh sentinel practices (as of 06/06/2021).



* The Moving Epidemic Method (MEM) threshold calculated for Wales ILI consultation rates is 11.1 per 100,000. MEM thresholds used in this chart are based on influenza from 2010-11 to 2018-19 seasons. Caution should be used when comparing consultation rates from March 2020 onwards to previous periods due to the changes in health-seeking behaviours brought about by the COVID-19 pandemic.

Figure 2. Clinical consultation rate for ILI per 100,000 practice population in Welsh sentinel practices (week 48 1996 – week 22 2021).



* Reporting changed to Audit+ surveillance system


Table 1. Age-specific consultations (per 100,000) for ILI in Welsh sentinel practices, week 17– week 22 2021 (as of 30/05/2021).

Age group	17	18	19	20	21	22
< 1	0.0	0.0	0.0	0.0	0.0	0.0
1 - 4	0.0	0.0	0.0	0.0	0.0	0.0
5 - 14	0.0	0.0	2.4	0.0	0.0	0.0
15 - 24	0.0	0.0	2.3	0.0	0.0	0.0
25 - 34	0.0	0.0	2.1	2.0	0.0	2.0
35 - 44	0.0	0.0	2.1	2.0	0.0	2.1
45 - 64	0.0	0.0	0.0	0.0	0.0	0.0
65 - 74	0.0	2.2	0.0	0.0	0.0	0.0
75+	0.0	0.0	0.0	0.0	0.0	0.0
Total	0.0	0.3	1.0	0.5	0.0	0.5

Table 2. Age-specific consultations (per 100,000) for ARI in Welsh sentinel practices, week 17 – week 22 2021 (as of 06/06/2021).


Age group	17	18	19	20	21	22
< 1	333.1	524.1	509.1	304.1	634.2	367.3
1 - 4	302.4	395.0	326.6	271.3	380.2	455.4
5 - 14	78.7	65.9	82.2	58.5	57.2	36.7
15 - 24	75.9	54.8	61.7	45.3	44.3	84.7
25 - 34	40.2	51.2	64.5	38.4	55.5	53.7
35 - 44	55.8	52.8	33.4	60.5	51.1	36.9
45 - 64	40.6	51.9	56.7	40.7	35.5	39.6
65 - 74	32.9	42.9	69.6	49.9	34.6	30.7
75+	56.3	50.1	54.0	40.5	66.8	51.7
Total	63.3	69.0	72.8	57.1	63.8	64.1

Figure 3. Specimens submitted for virological testing by sentinel GPs as of 06/06/2021, by week of sample collection, week 23 2020 to week 22 2021.



* Tested negative for influenza, adenovirus, rhinovirus, RSV, parainfluenza, mycoplasma, human metapneumovirus, enterovirus, bocavirus and coronaviruses.

Figure 4. Specimens submitted for virological testing for hospital patients and non-sentinel GPs as of 06/06/2021 by week of sample collection, week 23 2020 to week 22 2021.



This chart summarises respiratory panel test data and does not include data for patients tested SOLEY for SARS-CoV2. Combined data for tests carried out in Public Health Wales Microbiology: Cardiff laboratory, provided by Public Health Wales Microbiology Cardiff Specialist Virology Centre. This chart summarises individual test results, patients who are positive for multiple infections within a given week will appear multiple times.

Figure 5. Specimens from hospital patients submitted for RSV, Influenza and SARS-CoV2 testing only, as of 06/06/2021 by week of sample collection, week 46 2020 to week 22 2021.






Figure 6. Specimens submitted for virological testing for ICU patients, by week of sample collection, week 23 2020 to week 22 2021.




This chart summarises respiratory panel test data and does not include data for patients tested SOLEY for SARS-CoV2.

Figure 7. RSV incidence rate per 100,000 population aged under five years, week 31 2017 to week 22 2021.



Calls to NHS Direct Wales

Figure 8. Influenza related calls to NHS Direct Wales¹ (as a percentage of total calls) from week 23 2017 - week 22 2021 (as of 06/06/2021).



¹ Data supplied by Health Statistics and Analysis Unit, Welsh Government.

Flu related calls are the sum of calls recorded as 'cold/flu', 'cough', 'headache', 'fever' and 'sore throat'. Following changes to the NHS Direct calls system, including the start of the 111 pilot, there has been a change in the way in which denominator data are calculated for this chart, NHS Direct Wales now count the total number of nurse triaged calls (ie calls which could have symptom data recorded against them), note that 111 includes out-of-hours calls.

Influenza Vaccine Uptake in Wales

Table 3. Uptake of influenza immunisations in GP Practice patients, school children and NHS staff in Wales 2020/21 (as of 23/03/2021).

Influenza immunisation uptake in the 2020/21 season	
People aged 65y and older	76.5%
People younger than 65y in a clinical risk group	51.0%
Children aged two & three years	56.3%
Children aged four to ten years*	72.4%
NHS staff	63.4%
NHS staff who have direct patient contact	65.2%

* In school sessions carried out so far.

The end of season report Influenza in Wales 2019/20 is available to download and contains a full breakdown of vaccination uptake amongst eligible groups.

Link to report: <http://www.wales.nhs.uk/sites3/page.cfm?orgid=457&pid=55714>

Influenza activity – UK and international summary

- As of week 21, the majority of community and syndromic indicators remained low in the UK. GP ILI consultations remained stable in Northern Ireland at 0.4 per 100,000 and in Scotland at 0.2 per 100,000, well below the baseline intensity thresholds for both countries. The weekly ILI GP consultation rate in England reported through the RCGP system decreased to 0.7 per 100,000, still well below the MEM threshold for baseline activity (12.2 per 100,000).
- During week 21, none of the 2,195 respiratory test results reported through Public Health England's DataMart scheme tested positive for influenza. UK summary data are available from the [Public Health England National Influenza and COVID-19 Surveillance Report](#).
- The WHO and the European Centre for Disease Prevention and Control (ECDC) reported that as of week 21, influenza activity remained at inter-seasonal levels across the WHO European Region.

Source: Flu News Europe: <http://www.flunewseurope.org/>

- The WHO reported on 07/06/2021 that globally, despite continued or even increased testing in some countries, influenza activity remained at lower levels than expected for this time of year. In the temperate zone of the northern hemisphere, influenza activity remained below baseline, though detections of influenza B-Victoria lineage slightly increased, especially in China. In the temperate zones of the southern hemisphere, influenza activity remained at an inter-seasonal level. In the Caribbean and Central American countries, there were no influenza detections reported. In tropical South America, no influenza detections were reported. In tropical Africa, influenza detections were reported in some countries in Western and Middle Africa. In Southern Asia, influenza detections were reported in India. In South East Asia, no influenza detections were reported. Worldwide, influenza B accounted for the majority of the very low numbers of detections reported.
- Based on FluNet reporting (as of 04/06/2021), during the time period from 10/05/2021 – 23/05/2021, National Influenza Centres and other national influenza laboratories from 89 countries, areas or territories reported influenza surveillance data. The WHO Global Influenza Surveillance and Response System laboratories tested more than 284,780 specimens during that time period, 764 were positive for influenza viruses, of which 69 were typed as influenza A (of the subtyped influenza A viruses, 11 were influenza A(H1N1)pdm09 and nine were influenza A(H3N2) and 695 influenza B (of the characterised influenza B viruses four belonged to the B-Yamagata lineage and 642 to the B-Victoria lineage).

Source: WHO influenza update:

<https://www.who.int/teams/global-influenza-programme/surveillance-and-monitoring/influenza-updates/current-influenza-update>

Australia and New Zealand update

- In New Zealand, during the week ending 28/05/2021, the GP influenza-like illness activity (ILI) consultation rates overall have been lower than the weekly rates seen at this time in 2020, and are following a similar trend to historical rates. There have been no influenza viruses detected this season to date, rhinovirus has been the most commonly detected virus but numbers are low.
- In Australia, according to the latest available update (fortnight ending 23/05/2021), influenza-like illness (ILI) activity in the year to date remains at low levels. To date, the majority of nationally reported laboratory-confirmed influenza cases were influenza A (73.1%).

Source: Australian Influenza Surveillance Report and Activity Updates.

<https://www1.health.gov.au/internet/main/publishing.nsf/Content/cda-surveil-ozflu-flucurr.htm#current>

Respiratory syncytial virus (RSV) in North America

- The USA CDC has reported an out of season increase in RSV activity, with an increase in sample positivity since early March 2021.

Source: CDC RSV national trends

<https://www.cdc.gov/surveillance/nrevss/rsv/natl-trend.html>

Coronavirus disease 2019 (COVID-19) – UK and international summary

- The number of confirmed cases in Wales reported as at 09/06/2021 is 213,411, with 223 newly reported in the previous 24 hours. The cumulative number of suspected COVID-19 deaths in confirmed cases in hospitals and care homes reported to Public Health Wales is 5,570, with no new deaths reported in the previous 24 hours. The cumulative number of registered deaths in Welsh residents where COVID-19 was mentioned in the death certificate as at 2021 week 20 was 7,880. Latest COVID-19 data from Public Health Wales is available from: <https://phw.nhs.wales/topics/latest-information-on-novel-coronavirus-covid-19/> Public Health Wales produce a daily statement on COVID-19, available from: <https://covid19-phwstatement.nhs.wales/>
- As at 08/06/2021, there have been 4,528,442 reported confirmed cases of COVID-19 in the UK, of which 6,048 were newly reported in the previous 24 hours. The total deaths within 28 days of a positive test was 127,854, with 13 reported in the previous 24 hours. Latest UK data is available from: <https://coronavirus.data.gov.uk/>
- As at 09/06/2021, WHO have reported 173,609,772 confirmed COVID-19 cases globally, with 275,173 reported in the previous 24 hours. There have been 3,742,653 deaths, of which 7,016 were reported in the previous 24 hours. Daily WHO situation updates are available from: <https://covid19.who.int/>

Middle East respiratory syndrome coronavirus (MERS-CoV) – latest update from WHO and ECDC

- On 14/04/2021 WHO reported seven additional cases of Middle East Respiratory Syndrome coronavirus (MERS-CoV), including three associated deaths. Globally, 2,574 laboratory confirmed cases of human infection with MERS-CoV, including 886 associated deaths, have officially been reported to WHO since 2012.
Source: WHO Global Alert and Response website: <https://www.who.int/emergencies/disease-outbreak-news>
- The majority of the MERS cases continue to be reported from the Middle East, and specifically from Saudi Arabia. Rapid risk assessments of the situation from ECDC, which contain epidemiological updates and advice for travellers and healthcare workers, are available from: <https://ecdc.europa.eu/en/middle-east-respiratory-syndrome-coronavirus>
- Further updates and advice for healthcare workers and travellers are available from WHO: <http://www.who.int/emergencies/mers-cov/en/> and from NaTHNaC: <https://travelhealthpro.org.uk/news/237/mers-cov-update-travelhealthpro-country-pages>

Human infection with avian influenza A(H7N9), China

- Since February 2013, a total of 1,568 laboratory-confirmed cases of human infection with avian influenza A(H7N9), including at least 616 deaths, have been reported: http://www.fao.org/ag/againfo/programmes/en/empres/H7N9/Situation_update.html
- The risk of international spread of avian influenza A(H7N9) is considered to be low at present. However, it is important that clinicians are aware of the possibility of human infection with animal influenza, in persons presenting with severe acute respiratory disease, while travelling or soon after returning from an area where avian influenza is a concern. WHO Global Alert & Response updates: <https://www.who.int/emergencies/disease-outbreak-news>

Links:

Public Health Wales influenza surveillance webpage:

<http://www.wales.nhs.uk/sites3/page.cfm?orgid=457&pid=25480>

Public Health Wales COVID-19 data dashboard:

<https://public.tableau.com/profile/public.health.wales.health.protection#!/vizhome/RapidCOVID-19virology-Public/Headlinesummary>

GP Sentinel Surveillance of Infections Scheme:

<http://www.wales.nhs.uk/sites3/page.cfm?orgid=457&pid=27918>

NICE influenza antiviral usage guidance:

<http://www.nice.org.uk/Guidance/TA158>

Wales influenza information:

<https://phw.nhs.wales/topics/flu/>

England influenza and COVID-19 surveillance:

<https://www.gov.uk/government/statistics/national-flu-and-covid-19-surveillance-reports>

Scotland seasonal respiratory surveillance:

<https://beta.isdscotland.org/find-publications-and-data/population-health/covid-19/weekly-national-seasonal-respiratory-report/>

Northern Ireland influenza surveillance:

<https://www.publichealth.hscni.net/directorate-public-health/health-protection/seasonal-influenza>

European Centre for Communicable Disease:

<http://ecdc.europa.eu/>

European influenza information:

<http://flunewseurope.org/>

Advice on influenza immunisation (for NHS Wales users)

<http://nww.immunisation.wales.nhs.uk/home>

For further information on this report, please email Public Health Wales using:

surveillance.requests@wales.nhs.uk